## POSITIONS AND AREAS OF SUN SPOTS-Continued

	ì			Heli	ograph	ic				
Date	East- ern stand- ard time	Mount Wilson group No.	Dif- fer- ence in longi- tude	Lon- gi- tude	Lati- tude	Dis- tance from cen- ter of disk	Area of spot or group	Spot count	Plate qual- ity	Observatory
1939 Apr. 29	h m 10 41	6418 6417 6414 6413 6415 6410 6410 6412 6406 6407 6408 6405 6409	-76 -42 -39 -34 -31 -7 +81 +41 +42 +47 +55 +85	117 151 154 159 162 186 201 224 235 240 248 278 (193)	+11 -11 -20 +21 +12 +31 -11 -15 +30 +5 +23 -13 (-4)	77 42 41 41 35 35 11 32 42 53 48 60 84	388 97 48 194 24 24 242 48 776 630 97 145 242 2,955	7 12 9 25 3 8 30 15 30 20 1 3 7	VG	Mount Wilson.
Apr. 30	9 0	6418 6419 6414 6417 6414 6413 6415 6410 6412 6407 6406 6408	-63 -43 -38 -29 -21 -19 +14 +54 +60 +68	118 138 143 152 153 160 162 200 225 235 241 249 (181)	+12 -15 -21 -21 +20 +11 -11 +30 -15 +44 +22 (-4)	64 440 300 333 311 244 211 4461 6170	776 48 97 121 61 170 48 145 630 485 97 73 2,799	15 9 10 15 15 30 5 12 4 16 21 1	VG	Do.

Mean daily area for 30 days=2,133.

\*Not numbered.
Plate quality=F=fair; G=good; VG=very good; P=poor.

## PROVISIONAL SUNSPOT RELATIVE NUMBERS FOR APRIL 1939

[Dependent alone on observations at Zurich]

Data furnished through the courtesy of Prof. W. Brunner, Eidgen. Sternwarte, Zurich. Switzerlandl

April 1939	Relative numbers	April 1939	Relative numbers	April 1939	Relative numbers
1	Ec 34	11	a 103	21	125
3	Mc 85 83	12 13	a 100 EWcc 112	22 23	a 115 Macd 152
4	a 82 74	14 15	Eaac 126 b 121	24 25	151 a 134
	,-				
6	ad 70 Ecd 63	16 17	d 141 109	26 27	abd 134
8	dd 89	18 19	102 ad 94	28 29	ad
10	Wc 98	20	Macdd 125	30	140

Mean, 26 days = 106.2

Middle, large bright chromospheric eruption

	n.	774	4	7/4	
April 16	8	33 (	to 9	45 M.	
April 16	8	35	9	50 W.	
April 21	9	00	9	20 E.	
April 22	8	25	8	45 M.	
April 24	11	15	11	30 W.	

a = Passage of an average-sized group through the central meridian. b = Passage of a large group through the central meridian. c = New formation of a group developing into a middle-sized or large center of activity: E. on the eastern part of the sun's disk; W, on the western part; M, in the central-circle

d = Entrance of a large or average-sized center of activity on the east limb.

## AEROLOGICAL OBSERVATIONS

[Aerological Division, D. M. LITTLE in charge]

By B. Francis Dashiell

The 362 airplane and 233 radiosonde upper-air observations shown in tables 1 and 1a for the month of April also include the first of a series of radiosonde reports from Bermuda. Of all observations made exclusively within the United States, 96 percent and 85 percent reached 4 and 5 kilometers, respectively. Radiosonde observations showed some improvement, with 97, 92, 73, and 38 percent of all flights reaching 5, 10, 15, and 18 kilometers, respectively, while a few individual ascents rose to 23 kilometers. At Oakland, Calif., and Washington, D. C., 50 percent and 47 percent, respectively, of all flights launched at the surface attained 18 kilometers. The April wind resultants include those from 3 new pilot balloon stations operating at Des Moines, Iowa, Milwaukee, Wis., and Pueblo, Colo. A detailed explanation of tables 1, 1a, 2, 3, and 4, and charts VIII, IX, X, XI, and XII, will be found in the January 1939 issue of the Monthly Weather Review.

The weather in April was in contrast with that which prevailed during March over the eastern half of the country. As shown on chart I, mean surface temperatures (° F.) were subnormal east of the Mississippi Valley with the coldest weather over the Great Lakes region, and abnormal temperatures occurred over Nevada and the interior of California. Above the surface, in the free air, the mean temperatures (° C.) for April were lowest over Sault Ste. Marie, Mich., and Fargo, N. Dak., at all levels up to 8 kilometers; over Fargo, N. Dak., from 9 to 12 kilometers; over Oklahoma City, Okla., from 13 to 18 kilometers; and over Oklahoma City, Okla., and Oakland, Calif., at 19 and 20 kilometers. However, the lowest mean temperatures for the current month were recorded over Bermuda, between 12 and 18 kilometers.

A minimum upper-air mean temperature of -69.5° C. was reported from Bermuda, while the lowest for the United States (-63.2° C.) was recorded at Oklahoma City, Okla.; both occurring at 17 kilometers. Highest mean temperatures for the month were recorded over Pensacola, Fla., at 0.5, 1, and 5 kilometers; over El Paso, Tex., at all levels from 1.5 to 4 kilometers; at Oakland, Calif., from 6 to 9 kilometers; over Nashville, Tenn., at 10 and 11 kilometers; and over Sault Ste. Marie, Mich., from 12 to 20 kilometers, inclusive.

The April mean free-air temperatures listed in tables 1 and 1a were seasonally higher at all levels up to 5 kilometers than during March. At the 5-kilometer level the current month was cooler than April 1938, except over stations in the far Northwest and in California. Above 5 kilometers at all stations April was warmer than the preceding month, but Oakland, Calif., became cooler in

the levels higher than 10 kilometers.

A center of low mean pressure was indefinitely located north of the Great Lakes region and northeast toward Newfoundland, as shown on charts VIII, IX, X, and XI. At these same levels high pressure prevailed over Bermuda and extended westward in a belt to Pensacola, Fla., El Paso, Tex., and San Diego, Calif. Pressures during April were slightly higher in the North and lower in the South than in March. The pressure differences between the "high" and "low" areas, or gradient between Sault Ste. Marie, Mich., and Pensacola, Fla., at each level, increased with altitude up to 5 kilometers, but were found to be less than the differences noted in March.

Mean relative humidity, from the surface up to 8 kilometers, was highest over Sault Ste. Marie, Mich., and

comparatively high over Washington, D. C., at all levels above 4 kilometers. Bermuda reported high humidity at 0.5, 1, 1.5, and 2 kilometers, but at the higher levels the air was much drier. The driest air in the lower levels appeared over El Paso, Tex., and Norfolk, Va.; and over Oakland, Calif., Oklahoma City, Okla., and Bermuda, at all levels above 5 kilometers.

The April resultant winds, based on 5 a. m. and 5 p. m. observations, as shown on charts VIII, IX, X, and XI, showed northwesterly directions at 1.5 kilometers except for an area that extended from Texas to the Atlantic, and another in the far Northwest. At 3, 4, and 5 kilometers the directions were northwesterly except over the extreme southeastern portion of the United States, and Cuba and Mexico. At the four levels shown by the charts, winds from the northwest quadrant prevailed in 55, 79, 83, and 87 percent of all cases at 1.5, 3, 4, and 5 kilometers respectively. Even at the highest levels some southwesterly winds persisted. The April resultant wind directions were slightly north of those recorded in March, and this was particularly noticeable over the eastern half of the country at 3 kilometers.

Comparing the April 5 a. m. (E. S. T.) resultant wind directions at 1.5 and 3 kilometers (charts VIII and IX) with normal directions computed for the same levels at 24 well-distributed stations, it was found that most of the current winds departed from the normal directions by a clockwise rotation. However, outstanding exceptions occurred at Key West, Fla., and Atlanta, Ga., where the departures from normal were oriented by a counterclockwise rotation. Medford, Oreg., and Oakland, Calif., showed the largest degree of clockwise departures at 1.5 kilometers. Using these 5 a. m. normals in comparison with 5 p. m. observations at 4 and 5 kilometers (charts X and XI), it was found that large counterclockwise departures existed over Key West, Fla., and Medford, Oreg. Large clockwise departures were noted over Salt Lake City, Utah, at 3 and 4 kilometers. Resultant velocities for April were greater than normal at most of the stations for which normals have been computed, particularly over the Central States. The highest departure noted was +9.4 m. p. s. over Seattle, Wash., at 5 kilometers.

Table 2 shows 5 p. m. (E. S. T.) resultant wind directions and velocities at a number of selected stations at Comparing 20 of these stations with their 5 all levels. a. m. normals, it was found that clockwise departures occurred at all levels over Cheyenne, Wyo., and counter-clockwise departures at Sault Ste. Marie, Mich., and Washington, D. C. The most outstanding departures were noted at Salt Lake City, Utah, Omaha, Nebr., Med-ford, Oreg., Seattle, Wash., and Fargo, N. Dak.; while the winds at Chicago, Ill., Oakland, Calif., and St. Louis, Mo., were about normal Resultant valocities were higher were about normal. Resultant velocities were higher than normal in most cases, except at Cheyenne, Wyo., and San Diego, and Oakland, Calif. There were excessive velocities at Omaha, Nebr., and Seattle, Wash., being as much as +8.4 m. p. s. over the latter station at 5 kilometers.

The resultant winds for April, as shown in table 2, when compared to the winds of the previous month, showed higher velocities during April at most stations with the exception of Winslow, Ariz., Las Vegas, Nev., Havre, Mont., Fargo, N. Dak., Billings, Mont., Abilene, Tex., Albuquerque, N. Mex., Oklahoma City, Okla., St. Louis, Mo., and Chicago, Ill. The April resultant velocities at Medford, Oreg., and Miami, Fla., were considerably higher at all levels (up to 12 kilometers at the former) than during March. At Chicago, Ill., Fargo, N. Dak., Oklahoma City, Okla., St. Louis, Mo., and Winslow, Ariz, the March directions generally were south of (counterclockwise rotation) the April directions, and the wind velocities were higher.

The resultant winds for April at 1.5 and 3 kilometers (table 2) based on 5 p. m. observations, when compared with the current winds (charts VIII and IX) based on 5 a. m. observations, showed that the morning directions were definitely more northerly than the 5 p. m. directions. The 5 a.m. velocities, too, were slightly higher in most cases. However, at Miami, Fla., the 5 a. m. resultant directions were more southerly than at 5 p. m. at both levels by an average of  $28^{\circ}$ , and the resultant velocities lower by an average of -1.8 m. p. s.

The maximum altitudes reached by pilot balloons were 17,757 meters at Denver, Colo., and 17,556 meters at Omaha, Nebr. Altitudes over 15 kilometers were reached at 13 stations; from 10 to 15 kilometers at 39 stations, from 5 to 10 kilometers by 31 stations; and only one station failed to reach 5 kilometers. Maximum altitudes during April were somewhat higher than those reached in March, being confined to the Great Lakes region, Ohio Valley, New England, Texas, the southern Rocky Mountain region and southern Pacific coast. Practically all maximum altitudes were reached during the closing days of April-on the 28th over the northern Rocky Mountains, on the 29th over the north-central States and Upper Mississippi Valley, and on the 30th over the lower Ohio and Mississippi Valleys.

Table 3 shows the maximum wind velocities that occurred over the United States during April. Some relatively high wind speeds were recorded between the surface and 2.5 kilometers, the greatest being 51.4 m. p. s. from the north on the 5th, over Sandberg, Calif. This station also reported a maximum wind between 2.5 and 5 kilometers (56.1 m. p. s.) on the 5th. Above 5 kilometers, a maximum velocity of 80 m. p. s. from the northwest (179 miles per hour) occurred over Redding, Calif., on the 4th, at 13,270 meters. These velocities appear to be the highest recorded in the upper air over California in the levels below 2.5 kilometers, and above 5 kilometers.

The mean insentropic chart <sup>1</sup> for April shows two moist

tongues: One enters the country in Texas and curves anticyclonically to the northeast and east. This tongue was present on about one-third of the days, which is not characterized by winds blowing along its axis. cipitation excess above normal in the Northeastern States seems to be associated with this moist tongue. Another moist tongue was off the Atlantic coast, also on about one-third of the days; its northern portion was located by means of 22 daily radiosondes released by the International Ice Patrol cutters S. S. Champlain and S. S. Chelan at about latitude 44.5° N., longitude 48.5° W. In the west, the effect of the plateau is seen in the northward bulge of the contour and moisture lines.

I This chart and the following discussion have been prepared by the Air Mass Section of the Meteorological Research Division.

Table 1.—Mean free-air barometric pressures (P.) in mb., temperatures (T.) in °C., and relative humidities (R. H.) in percent obtained by airplanes during April 1939

								_				A	ltitu	de (n	neters)	m, s	. 1.										f .	
Thatian and along time in materia		Surfa	се			500			1,000	i		1,500			2,000			2,500			3,000			4,000			δ,000	
Stations and elevations in meters above sea level	Num- ber of obser- va- tions	Р.	т.	R. H.	P.	т.	R. H.	P.	т.	R. H.	P.	т.	R. H.	Р.	т.	R. H.	P.	т.	R. H.	P.	т.	R. H.	P.	т.	R. H.	P.	T.	R
Billings, Mont. (1,090 m.) Cheyenne, Wyo. (1,873 m.) Chicago, Ill. (187 m.) Coco Solo, C. Z.! (15 m.) El Paso, Tex. (1,193 m.) Lakehurst, N. J.! (39 m.) Norfolk, Va.! (10 m.) Pearl Harbor, T. H.! (6 m.) Pensacola, Fla.! (13 m.) Est. Thomas, V. I.! (8 m.) Salt Lake City, Utah (1,288 m.)	30	892 810 991 1, 012 881 1, 011 1, 017 1, 016 1, 017	7.1	86 36 74 76 80 89	954 958 955 959 960 960	11. 4 19. 1 16. 0	94 67 47 76 75	905 898 903 906 905	2, 9 8, 5 15, 9 14, 7	67 46 78 67	854 850 844 849 853 853	1. 2 17. 2 15. 6 0. 6 5. 1 13. 4 12. 0	32 69 50 76 65	797 792 805 800 793 799 804 803	3.8 4.5 -1.2 15.1 13.0 -1.8 2.8 11.2 9.7	53 62 60 70 31 69 46 67 56	750 750 744 759 754 744 751 756 756	0. 2 2. 9 -3. 4 13. 2 10. 0 -3. 8 0. 7 8. 9 7. 6	30 69 40 61 47	698 715 710 699 705 712 711	-0.6 -6.0 10.6 6.4 -5.9 -2.2 6.9 4.6	53 50 49 31 67 39 54 45	634 627 614 621 630 628	-1.1 -11.1 -6.8 2.1 -1.4	55 48 39 33 62 29 41 35	545 538 552 546 554	-13.3 -7.7	3 3
an Diego, Calif.! (10 m.) eattle, Wash.! (10 m.) pokane, Wash. (597 m.)	1 27	1,016 1,019	14. 4 10. 7 6. 3	80 67 69	958	13. 3 8. 1	77 66	904 904 902	14.7 5.5 9.4	65	851	3, 2	52 50 61 50	800 802 800 799	11.7 0.7 2.8	41 55 52	752 755 751 751	8.7 -1.2 -0.7	36 49 54	710 706 705	5. 2 -3. 7	34 47 52	627 620	-1.3 -9.2 -9.7	31 40	552	-9.3 -15.5	3

Table 1a.—Mean free-air barometric pressures (P) in mb., temperatures (T) in °C., and relative humidities (R H) in percent obtained by radiosonde during April 1939

												Sta	tions	and o	elevatio	ons i	n me	ters a	above	sea le	vel											
		Berm (50 1			Fa		N. Da 4 m.)	k.	Na		lle, Tei 0 m.)	nn.	o		d, Cali m.)	if.	Ole O	laho kla.	ma Ci (391 m	ty, .)	C	mah: (30)	a, Neb 0 m.)	r.	Sat	ult S fich.	te. Ma (221 n	rie, 1.)	Was	shing (13	ton, D	. C.1
Surface	Number of observations	Pressure	Temperature	Relative humidity	Number of observations	Pressure	Temperature	Relative humidity	Number of obser- vations	Pressure	Temperature	Relative humidity	Number of observations	Pressure	Temperature	Relative humidity	Number of tobservations	Pressure	Temperaturo	Relative humidity	Number of observations	Pressure	Temperature	Relative humidity	Number of observations	Pressure	Temperature	Relative humidity	Number of observations	Pressure	Temperature	Relative humidity
	24 24 24 24 24 24 24 21 18 13	964 909 856 805 758 712 629 554 486 425 370 276 237 202 172 1746 124 105 88	17. 9 16. 0 12. 4 9. 5 7. 0 5. 3 -2. 9 -9. 1 -15. 6 -22. 8 -30. 5 -38. 4 -60. 2 -46. 3 -54. 1 -60. 2 -63. 8 -64. 5 -69. 5 -66. 4 -69. 5	69 58 52 43 40 39 38 37 36	30 30 30 30 30 30 30 30 30 30	954 897 842 791 743 696 612 535 467 406 351 302 259 221 188 160 136 136 198 83	-3. 2 -5. 2 -7. 9 -13. 8 -20. 0 -26. 8 -34. 0 -41. 6 -48. 3 -54. 3 -59. 3 -59. 3 -58. 5	66 63 62 61 57 55 54 54	300 300 300 300 300 285 227 225 225 225 225 225 225 225 225 22	749 704 620 545 478 417 362 314 270 232 199 170 144 123 105	-1.3 -5.8 -11.8 -18.2 -25.2 -32.1 -39.4 -46.2 -51.9 -57.8 -57.8 -57.8	70 69 65 63 61 63 59 55 54 52 52	30 30 30 30 30 30 30 30 30 30	851 800 753 708 625 550 482 421 366 317 274 235 201 171 146 124 106 90 65	10. 4 11. 9 10. 4. 8 7. 7 7 4. 8 2. 2 2 -3. 6 -9. 9 -17. 1 -31. 7 -39. 3 -46. 5 -60. 4 -59. 9 -60. 6 -60. 7	31 30	28 27 26	480 419 365 315 272 233 199 169 144 123 104 89 75 64	11. 3 12. 1 11. 8 9. 3 6. 8 4. 8 4. 8 -4. 2 -11. 2 -25. 8 -33. 1 -40. 4 -54. 1 -59. 0 -61. 3 -63. 2 -63. 1 -62. 3	53 50 48 46 40 39 40 39 37	30 30 30 30 30 30 29 29 26 26	845 795 746 701 617 541 473 412 357 308 265 227 194 165 140 102 86	-58.6 $-59.3$	59 58 57 53 50 47 46 45	30 30 30 30 30 30 30 29 28 28 28 28 27 26	952 894 839 787 738 692 606 606 530 462 400 346 298 255 219 160 137 117 100 85 72	-1. 2 -2. 1 -4. 3 -6. 6 -8. 9 -11. 4 -16. 7 -23. 2 -29. 7 -36. 3 -42. 6 -54. 3 -54. 7 -54. 8 -55. 6 -57. 3	83 78 77 76 72 69 65 63 60 59 60	30 30 30 30 30 30 30 30 30	473 412 358 310 267 229 196 167 143 122 103 88 74	-56.0 -57.7	65 68 70 67 66 65 64 59 57 57 57 56

<sup>1</sup> Navy.
2 Flights discontinued temporarily.

Observations taken about 4 a. m. 75th meridian time, except by Navy stations along the Pacific coast and Hawaii where they are taken at dawn. Note.—None of the means included in this table are based on less than 15 surface or 5 standard-level observations.

<sup>&</sup>lt;sup>1</sup> Navy.
<sup>2</sup> Operated by Massachusetts Institute of Technology.

Observations taken about 4 a. m. 75th meridian time, except by Navy stations along the Pacific coast and Hawaii where they are taken at dawn.

Note.—None of the means included in this table are based on less than 15 surface or 5 standard-level observations.

Number of observations refers to pressure only as temperature and humidity data are missing for some observations at certain levels, also, the humidity data are not used in daily observations when the temperature is below -40° C.

## MONTHLY WEATHER REVIEW

Table 2.—Free-air resultant winds based on pilot-balloon observations made near 5 p. m. (E. S. T.) during April 1939 [Directions given in degrees from North (N=360°, E=90°, S=180°, W=270°)—Velocities in meters per second (superior figures indicate number of observations)]

<del></del>		<u> </u>		- 1		<u> </u>						·			<del></del>		1						<del></del>			
Altitude	Abii To (537	ex.	Albu qı N. 1 (1,55	ie, Mex.	Atla G (302		Billi Mc (1,09	nt.	Bo Ids (850	aho	N.	klyn, Y. m.)	Brovil Vil (7 1	le,	N.	falo, Y. m.)	ton,	ling- Vt. m.)	ton,	rles- S. C. m.)	W	enne, yo. 3 m.)	l	cago, ll. !m.)	na O	ncin- ati, hio ' m.)
(meters) m. s. l.	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity
Surface	287 213 241 268 277 278 284 291 291 293	0. 430 0. 520 1. 428 2. 727 4. 624 6. 923 9. 738 12. 920 14. 317 16. 815 16. 310	279 287	1. 5 <sup>30</sup> 2. 7 <sup>30</sup> 3. 3 <sup>30</sup> 4. 2 <sup>30</sup> 7. 8 <sup>36</sup> 12. 3 <sup>34</sup> 16. 0 <sup>22</sup> 20. 5 <sup>15</sup> 22. 8 <sup>13</sup>	260 258 258 252 260 277 270 281 289	9.324	314 307 311 297 285 281 295 293	2. 230 2. 629 4. 229 5. 427 7. 023 7. 518 8. 712	306 297 300 307 299 286 293	2. 130 3. 130 3. 230 3. 630 4. 130 5. 229 7. 322 7. 416	256 260 264 274 280 287	4. 0 <sup>27</sup> 6. 3 <sup>27</sup> 8. 0 <sup>26</sup> 10. 2 <sup>26</sup> 12. 7 <sup>22</sup> 15. 1 <sup>18</sup>	° 128 135 169 193 275 263 289 276 282	5. 029 4. 929 2. 925 1. 623 2. 220 3. 710 6. 618 10. 918 13. 010	251 264 258 258 268 269	4. 528 6. 928 7. 724 8. 617 8. 715 9. 912	299 276 273 263 274 285 293	1. 524 2. 623 3. 922 4. 621 5. 414 7. 316 9. 012	263 266	10, 426	292  293 299 293 291 306	4. 427 6. 427 7. 876 9. 023 9. 213	262 258 268 271 283 290 306	2. 4 <sup>27</sup> 3. 6 <sup>27</sup> 4. 6 <sup>23</sup> 7. 0 <sup>20</sup> 9. 313 10. 313 11. 011	254 269 295	3. 038 5. 729 7. 029 7. 720 7. 818 10. 111
Altitude	Т	Paso, er. 16 m.)	N.	rgo, Dak. 3 m.)	bo N.	ens- oro, . C. l m.)	M	vre, ont. om.)	T	ex. m.)	8.1	ron, Dak. Sm.)	N-	/egas, ev. m.)	Ro	ttle ock, rk. m.)	Or	lford, reg. m.)	F	ami, la. m.)	ap M	nne- olis, inn. m.)	Te	nville, nn. m.)	Orle L	ew eans, .a. m.)
(meters) m. s. l.	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity
Surface		3. 130 3. 010 2. 938 4. 329 6. 924 8. 924 11. 420	346 354 347 331 325 324 323 320	1. 429 4. 229 3. 627 4. 326 5. 321 6. 019 6. 116 9. 012	242 251 250 254 267 275 274 268 292	4. 229 5. 829 7. 229 9. 739 13. 424 15. 920 18. 618	288 271 267 275 277 288 290 282	2. 2 <sup>28</sup> 3. 8 <sup>28</sup> 3. 8 <sup>28</sup> 4. 5 <sup>24</sup> 6. 11 <sup>9</sup> 7. 11 <sup>6</sup> 9. 6 <sup>11</sup> 9. 2 <sup>10</sup>	142 164 234 289 285 280 277 293 295		324 316	8. 524	96 	1. 030 0. 830 0. 430 2. 230 3. 130 2. 323 3. 723 4. 827 8. 428 10. 520 14. 717 14. 414	284	1. 0 <sup>29</sup> , 2. 3 <sup>29</sup> , 3. 4 <sup>29</sup> , 5. 7 <sup>29</sup> , 6. 9 <sup>28</sup> , 7. 4 <sup>22</sup> , 9. 0 <sup>20</sup> , 12. 8 <sup>16</sup> , 14. 3 <sup>12</sup>	318 315 298 266 238 237 244 251 258 260 272 255 240	2. 330 2. 650 2. 230 1. 530 2. 230 3. 429 4. 729 7. 197 8. 726 10. 524 11. 217 10. 413 12. 510	130 143 198 230 255 264 265 251		304 312 282 271 278 296 319	2, 526 2, 926 3, 226 5, 326 6, 422 6, 314 9, 710	250 229 237 250 256 263 275 272	2. 430 3. 330 4. 530 6. 130 7. 327 8. 328 10. 421 13. 219	262 189 216 247 262 278 282 290 304	0. 130 2. 430 2. 826 3. 345 5. 323 6. 523 7. 120 9. 016 12. 210
Altitude (meters)	Ci	land, alif. m.)	City.	homa Okla. m.)	Ne	aha, ebr. 3 m.)	N.	no, ev. 6 m.)	M	Louis, Io. m.)	City,	Lake Utah 1 m.)		Diego, lif. m.)	P.	Juan, R. m.)	Ma	t Ste. arie, ich. m.)	W	ttle, ash. m.)	Wε	kane, ish., m.)	Was ton, 1	hing- D. C. m.)		slow, riz. 8 m.)
m. s. l.	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Dírection	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity	Direction	Velocity
Surface	260 290 301 304 297 287 288 285 281 264	3. 728 2. 628 2. 638 2. 428 1. 828 2. 128 4. 626 7. 023 9. 022	258 275 270 277 277 282 289	14. 818	289 299 297	3. 928 5. 226	260 289 287	1. 830 1. 730 1. 630 2. 130 3. 329 4. 928 5. 826 8. 623 9. 917 8. 813	233 260 252 259 275 288 292 291	1. 5 <sup>28</sup> 2. 7 <sup>28</sup> 4. 5 <sup>26</sup> 5. 7 <sup>24</sup> 7. 8 <sup>20</sup> 8. 91 <sup>7</sup> 10. 2 <sup>16</sup> 15. 0 <sup>12</sup>	308 308 289 256 250 294 307	3. 0 <sup>30</sup> 3. 2 <sup>30</sup> 2. 5 <sup>30</sup> 3. 3 <sup>29</sup> 3. 8 <sup>28</sup> 3. 8 <sup>21</sup> 4. 113	279 293 316 339 16 349 347 321 294 291	4. 029 2. 429 2. 323 1. 123 1. 121 1. 620 2. 017 3. 916 6. 013 9. 311	112 118 128 157	6. 0 <sup>28</sup> 4. 7 <sup>18</sup> 3. 6 <sup>14</sup>	° 301 315 313 299 288 297 299 302	5. 020 8. 519	207 225 239 284	2. 426 2. 329 3. 227 4. 023 4. 521 5. 518 5. 817 10. 212 12. 910	263 278	3. 0 <sup>30</sup> 3. 0 <sup>30</sup> 3. 6 <sup>29</sup> 5. 4 <sup>36</sup> 5. 7 <sup>22</sup> 5. 5 <sup>19</sup> 6. 7 <sup>17</sup> 10. 2 <sup>14</sup>	268 273	1. 629 4. 629 7. 328 9. 526 10. 723 13. 322 16. 318	248 253 253 280 294 312 308 290	3. 330 3. 030 3. 430 3. 730 4. 928 6. 624 8. 523 14. 449 20. 318

Table 3 .- Maximum free-air wind velocities, (M. P. S.), for different sections of the United States based on pilot balloon observations during April 1939

		Surface	to 2,500	mete	ers (m. s. l.)		Between 2,	500 and	5 <b>,00</b> 0	meters (m. s. l.)		Above !	,000 me	ters	(m. s. l.)
Section	Maximum velocity	Direction	Altitude (m.) m. s. l.	Date	Station	Maximum velocity	Direction	Altitude (m.) m. s. l.	Date	Station	Maximum velocity	Direction	Altitude (m.) m.s.l.	Date	Station
Northeast 1 East-Central 2	42. 0 39. 1	WNW		3 10	Kylertown, Pa Nashville, Tenn	45. 6 48. 4	sw WNW	3, 590 4, 530	10 12	Columbus, Ohio Knoxville, Tenn	57. 0 49. 6	NW NNW		3 4	Cleveland, Ohio. Greensboro, N.
Southeast <sup>3</sup> North-Central <sup>4</sup> Central <sup>3</sup> South-Central <sup>6</sup> Northwest <sup>7</sup> West-Central <sup>8</sup> Southwest <sup>9</sup>	36. 8 43. 6 41. 9 40. 0 30. 1 38. 6 51. 4	SW	1, 970 2, 500	19 25 9 16 5 13 5	Charleston, S. C. Minneapolis, Minn. Evansville, Ind. Del Rio, Tex. Boise, Idaho. Modena, Utah. Sandberg, Calif.	40. 5 40. 6 52. 0 35. 0 39. 0 48. 0 56. 1	SWSWNWNWNWNWNW.	4, 670 2, 930 4, 310 4, 830	19 12 9 1 15 7 5	Charleston, S. C	43. 0 46. 1 52. 8 41. 4 45. 2 80. 0 55. 2	W	11, 550 11, 240 5, 210 17, 380 13, 270	23 21 9 7 28 4 10	Charleston, S. C. Huron, S. Dak. Wichita, Kans. Little Rock, Ark Billings, Mont. Redding, Calif. Albuquerque, Mex.

<sup>1</sup> Maine, Vermont, New Hampshire, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania, and northern Ohio.
2 Delaware, Maryland, Virginia, West Virginia, southern Ohio, Kentucky, eastern Tennessee, and North Carolina.
3 South Carolina, Georgia, Florida, and Alabama.
4 Michigan, Wisconsin, Minnesota, North Dakota, and South Dakota.
5 Indiana, Illinois, Iowa, Nebraska, Kansas, and Missouri.
6 Missssippi, Arkansas, Louisiana, Oklahoma, Texas (except El Paso), and western Tennessee.
7 Montana, Idaho, Washington, and Oregon.
8 Wyoming, Colorado, Utah, northern Nevada, and northern California.
9 Southern California, southern Nevada, Arizona, New Mexico, and extreme west Texas.

Table 4.—Mean altitudes and temperatures of significant points identifiable as tropopauses during April 1939, classified according to the potential temperatures (10-degree intervals between 290° and 399° A.) with which they are identified. (Based on radiosonde observations.)

Ì	Far	go, N. I	Oak.	Nasi	nville, T	enn.	Oak	dand, C	alif.	Oklaho	ma City	y, Okla.	On	aha, N	ebr.	Saul	t Ste. M Mich.		Wash	ington,	D. C.
	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture ° C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture ° C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture ° C.	Num- her of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture ° C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture ° C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture ° C.	Num- ber of cases	Mean alti- tude (km.) m. s. l.	Mean tem- pera- ture ° C.
290-299	5	7.9	-55.6														6.8	-46. 2			
300-309		8.0	-48.8				<b>-</b>			;-	7, 7	-46.0	<u>3</u>	7.8	-46.0	4	8.0	-50.7		5-5-	
310-319	15	9.6	-57.0		8. 5	-41.8	2	8. 1	-40.0	6	9. 1	-48.3	7	8.9	-49. 1	14	9.2	- 52. 6	3	9. 1 8. 5	-58.0
320-329	14	10. 9	-61.9	12	9. 6	-47.0	10	10.5	-52.5	10	9. 5	-46.8	17	10. 1	-53. 1	13	10. 2	-55.3	8	9.8	<b>-45.0</b>
330-339	6	11.7	-62.3	19	11.0	-54.5	20	11.4	-58.0	17	11.5	-58.6	17	11.5	-58.6	12	11. 2	-57.8	15	11.3	-49.5 -56.3
340-349	5	12. 5	-62.4	14	12.5	-60.6	22 13	13. 0	-65. 7	l io	12.6	-62.7	10	12. 4	-61.1	9	11.4	-56.3	10	12.4	-61.3
350-359	ĭ	13. 6	-67. O	1 78	12.8	-58.9	5	13. 1	-61.4	6	13. 6	-65. 2	7	12.6	-58. î	2	12.8	-60.5	7	13.6	-63. 6
360-369	ī	12. 2	-54.0	ľĭ	13. 1	-56.0	3	13, 3	-57. Ô	3	13. 5	-60.7	3	12.8	-55.7	4	12. 5	-56.0	l i	13.8	-69.0
370-379				5	14.2	-60.2	1	14.4	-61.0				3	13.8	-58.3	3	13. 1	-53. 7	l î	14. 2	-58.0
380-389	1	15.0	-64.0	2	14.8	59. 5	1	14. 2	-52.0	1	14.8	-58.0	4	14.3	-57.8				l		
390-399			l	1	15. 4	-60.0				1	16. 4	-70.0	1	15. 2	-58.0	1	13. 5	-48.0			_
All (weighted		ŀ	ļ	l						i							l	_			
means)		10.4	-58.8		11.6	54.9		11.8	-58. Q		11.5	-56.9		11.4	56. 0		10.2	-54.2		11.4	-56.5
Mean potential temperature	322.3				341.8			339.2	1		337.6	<u>.                                    </u>		338.5			328.6			337.2	<u></u>